## This Page Is Inserted by IFW Operations and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

			Ŷ.	
· · · · · · · · · · · · · · · · · · ·		·y·		
		•		
			A.C.	
	2	•		
·				
	. 9			•
			1.6	
				4
	•			
```				

## PATENT ABSTRACTS OF JAPAN

(11) Publication number:

10-060480

(43)Date of publication of application: 03.03.1998

(51) Int. CI.

C11D 3/37 C11D 10/02 C11D 17/08 //(C11D 10/02 C11D 1:72 C11D 1:722 C11D 3:37 C11D 3:20 )

(21) Application number: 08-218855

(71) Applicant: KAO CORP

(22) Date of filing:

20.08.1996

(72) Inventor: YAMADA ISAO

SHIGEMATSU KUNIYOSHI TERANISHI FUTOSHI

(54) LIQUID DETERGENT COMPOSITION

(57) Abstract:

PROBLEM TO BE SOLVED: To obtain a liquid detergent composition, having the moderate detergency for the so-called fancy wear and capable of further imparting excellent touch feeling and dimensional stability thereto by blending a specific aminomodified silicone derivative with plural surfactants.

SOLUTION: This liquid detergent composition comprises (A) 0.05-5wt.% amino-modified silicon derivative represented by formula I [(I) is 100-600; (m) and (n) are each a number so as to provide (I): (m)=(100:1) to (10:1); (m): (n)=(1:10) to (10:1); R is a 1-4C alkyI, etc.; R1 is a 1-4C alkyI; A is represented by formula II, etc., [(a) is 2-6; R2 is H or a 1-4C alkyI; (p) is 1-6; (q) is 1-20; (r) is 0-20; R3 is a 1-18C alkyI]; B is (CH2) a-0-(C2H40) x-(C3H60) y-R4 [(x) is 1-20; (y) is

0-20; R4 is a 1-10C alkyl] or R1] and (B) 5-70wt.% surfactant containing (i) 95-75wt.% nonionic

surfactant represented by formula III [R1 is an 8-18C alkyl, etc.; R2 is a 2-4C alkyl; R3 is H, methyl, etc.; (n) is the number of mol of an added alkylene oxide] and (ii) 5-15wt.% nonionic surfactant, represented by formula IV (R4 is R1; R5 is R3) and having 7-10 value of hydrophile-lipophile balance(HLB) and \*4wt.% content of a compound in which (n) is 0 and the number of mol of the compound having the highest content (nmax) satisfies formula V. The weight ratio of the ingredients A:B is (1:100) to (1:5).

RLOCK DO R

T

II.

T

IV.

 $\hat{\mathbf{y}}$ 

Rad (CHachan) e Re

Σ ြ\_\_\_\_\_\_ Υ, ≥ co%

LEGAL STATUS

[Date of request for examination]

07. 02. 1997

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

2863137

[Date of registration]

11. 12. 1998

[Number of appeal against examiner's decision of rejection]

[Date of requesting appeal against examiner's decision of rejection]
[Date of extinction of right]

Copyright (C); 1998, 2003 Japan Patent Office